arXiv:hep-ex/0308048v2 19 Mar 2004

Belle Preprint 2004-11
KEK Preprint 2003-143

Study of CP Violating Effects in Time Dependent $B^0(\bar{B}^0) \to D^{(*)\mp}\pi^{\pm}$ Decays

T. R. Sarangi,[7] K. Abe,[7] K. Abe,[38] T. Abe,[7] H. Aihara,[40] Y. Asano,[44] V. Aulchenko,[1] T. Aushev,[11] S. Bahinipati,[3] A. M. Bakich,[35] Y. Ban,[29] S. Banerjee,[36] U. Bitenc,[12] I. Bizjak,[12] S. Blyth,[23] A. Bondar,[1] M. Bračko,[18, 12] T. E. Browder,[6] M.-C. Chang,[23] P. Chang,[23] K.-F. Chen,[23] B. G. Cheon,[34] R. Chistov,[11] S.-K. Choi,[5] Y. Choi,[34] A. Chuvikov,[30] S. Cole,[35] M. Danilov,[11] M. Dash,[46] L. Y. Dong,[9] A. Drutskoy,[11] S. Eidelman,[1] V. Eiges,[11] S. Fratina,[12] N. Gabyshev,[7] A. Garmash,[30] T. Gershon,[7] G. Gokhroo,[36] J. Haba,[7] N. C. Hastings,[7] H. Hayashii,[21] M. Hazumi,[7] T. Higuchi,[7] L. Hinz,[16] T. Hokuue,[20] Y. Hoshi,[38] W.-S. Hou,[23] K. Inami,[20] A. Ishikawa,[7] H. Ishino,[41] R. Itoh,[7] H. Iwasaki,[7] M. Iwasaki,[40] J. H. Kang,[48] J. S. Kang,[14] P. Kapusta,[24] S. U. Kataoka,[21] H. Kawai,[2] T. Kawasaki,[26] H. Kichimi,[7] H. J. Kim,[48] H. O. Kim,[34] S. K. Kim,[33] T. H. Kim,[48] K. Kinoshita,[3] S. Korpar,[18, 12] P. Križan,[17, 12] P. Krokovny,[1] A. Kuzmin,[1] Y.-J. Kwon,[48] J. S. Lange,[4, 31] G. Leder,[10] S. H. Lee,[33] T. Lesiak,[24] J. Li,[32] S.-W. Lin,[23] D. Liventsev,[11] J. MacNaughton,[10] F. Mandl,[10] D. Marlow,[30] T. Matsumoto,[42] A. Matyja,[24] Y. Mikami,[39] W. Mitaroff,[10] K. Miyabayashi,[21] H. Miyake,[27] H. Miyata,[26] D. Mohapatra,[46] G. R. Moloney,[19] T. Nagamine,[39] Y. Nagasaka,[8] T. Nakadaira,[40] M. Nakao,[7] H. Nakazawa,[7] Z. Natkaniec,[24] S. Nishida,[7] O. Nitoh,[43] T. Nozaki,[7] S. Ogawa,[37] T. Ohshima,[20] T. Okabe,[20] S. Okuno,[13] S. L. Olsen,[6] W. Ostrowicz,[24] H. Ozaki,[7] P. Pakhlov,[11] H. Palka,[24] H. Park,[15] K. S. Park,[34] N. Parslow,[35] L. E. Piilonen,[46] M. Rozanska,[24] H. Sagawa,[7] S. Saitoh,[7] Y. Sakai,[7] M. Satapathy,[45] O. Schneider,[16] J. Schümann,[23] C. Schwanda,[10] A. J. Schwartz,[3] S. Semenov,[11] K. Senyo,[20] M. E. Sevior,[19] H. Shibuya,[37] V. Sidorov,[1] J. B. Singh,[28] N. Soni,[28] R. Stamen,[7] S. Stanič,[44, *] M. Starič,[12] K. Sumisawa,[27] T. Sumiyoshi,[42] S. Suzuki,[47] O. Tajima,[39] F. Takasaki,[7] M. Tanaka,[7] G. N. Taylor,[19] T. Tomura,[40] T. Tsuboyama,[7] S. Uehara,[7] T. Uglov,[11] K. Ueno,[23] Y. Unno,[2] S. Uno,[7] G. Varner,[6] K. E. Varvell,[35] C. H. Wang,[22] M.-Z. Wang,[23] M. Watanabe,[26] B. D. Yabsley,[46] Y. Yamada,[7] A. Yamaguchi,[39] Y. Yamashita,[25] J. Ying,[29] Y. Yusa,[39] J. Zhang,[7] Z. P. Zhang,[32] Y. Zheng,[6] V. Zhilich,[1] T. Ziegler,[30] and D. Žontar[17, 12]

(The Belle Collaboration)

[1]*Budker Institute of Nuclear Physics, Novosibirsk*
[2]*Chiba University, Chiba*
[3]*University of Cincinnati, Cincinnati, Ohio 45221*
[4]*University of Frankfurt, Frankfurt*
[5]*Gyeongsang National University, Chinju*
[6]*University of Hawaii, Honolulu, Hawaii 96822*
[7]*High Energy Accelerator Research Organization (KEK), Tsukuba*

[8]*Hiroshima Institute of Technology, Hiroshima*
[9]*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing*
[10]*Institute of High Energy Physics, Vienna*
[11]*Institute for Theoretical and Experimental Physics, Moscow*
[12]*J. Stefan Institute, Ljubljana*
[13]*Kanagawa University, Yokohama*
[14]*Korea University, Seoul*
[15]*Kyungpook National University, Taegu*
[16]*Swiss Federal Institute of Technology of Lausanne, EPFL, Lausanne*
[17]*University of Ljubljana, Ljubljana*
[18]*University of Maribor, Maribor*
[19]*University of Melbourne, Victoria*
[20]*Nagoya University, Nagoya*
[21]*Nara Women's University, Nara*
[22]*National United University, Miao Li*
[23]*Department of Physics, National Taiwan University, Taipei*
[24]*H. Niewodniczanski Institute of Nuclear Physics, Krakow*
[25]*Nihon Dental College, Niigata*
[26]*Niigata University, Niigata*
[27]*Osaka University, Osaka*
[28]*Panjab University, Chandigarh*
[29]*Peking University, Beijing*
[30]*Princeton University, Princeton, New Jersey 08545*
[31]*RIKEN BNL Research Center, Upton, New York 11973*
[32]*University of Science and Technology of China, Hefei*
[33]*Seoul National University, Seoul*
[34]*Sungkyunkwan University, Suwon*
[35]*University of Sydney, Sydney NSW*
[36]*Tata Institute of Fundamental Research, Bombay*
[37]*Toho University, Funabashi*
[38]*Tohoku Gakuin University, Tagajo*
[39]*Tohoku University, Sendai*
[40]*Department of Physics, University of Tokyo, Tokyo*
[41]*Tokyo Institute of Technology, Tokyo*
[42]*Tokyo Metropolitan University, Tokyo*
[43]*Tokyo University of Agriculture and Technology, Tokyo*
[44]*University of Tsukuba, Tsukuba*
[45]*Utkal University, Bhubaneswer*
[46]*Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061*
[47]*Yokkaichi University, Yokkaichi*
[48]*Yonsei University, Seoul*

Abstract

We report measurements of time dependent decay rates for $B^0(\bar{B}^0) \rightarrow D^{(*)\mp}\pi^{\pm}$ decays and extraction of CP violation parameters containing ϕ_3. Using fully reconstructed $D^{(*)}\pi$ events from a 140 fb^{-1} data sample collected at the $\Upsilon(4S)$ resonance, we obtain the CP violation parameters for $D^*\pi$ and $D\pi$ decays, $2R_{D^{(*)}\pi}\sin(2\phi_1 + \phi_3 \pm \delta_{D^{(*)}\pi})$, where $R_{D^{(*)}\pi}$ is the ratio of the magnitudes of the doubly-Cabibbo-suppressed and Cabibbo-favoured amplitudes, and $\delta_{D^{(*)}\pi}$ is the strong phase difference between them. Under the assumption of $\delta_{D^{(*)}\pi}$ being close to either 0 or 180°, we obtain $|2R_{D^*\pi}\sin(2\phi_1 + \phi_3)| = 0.060 \pm 0.040(\mathrm{stat}) \pm 0.019(\mathrm{sys})$ and $|2R_{D\pi}\sin(2\phi_1 + \phi_3)| = 0.061 \pm 0.037(\mathrm{stat}) \pm 0.018(\mathrm{sys})$.

PACS numbers: 12.15.Hh,13.25.Hw

The good agreement between direct measurements of $\sin 2\phi_1$ [1, 2] and the outcome of global fits to the CKM quark mixing matrix elements [3] strongly supports the standard model explanation of CP violation. To determine whether it is the complete description or whether additional factors come into play, further measurements of other CKM parameters are required. Among these parameters ϕ_3 is of particular importance. The measurements of time-dependent decay rates of $B^0(\bar{B}^0) \rightarrow D^{(*)\mp}\pi^{\pm}$ provide a theoretically clean method for extracting $\sin(2\phi_1 + \phi_3)$, since loop diagrams do not contribute to these decays [4].

There are two ways for a state which is initially B^0 to be found as $D^{(*)-}\pi^+$ at a later time t. It can occur either directly through a Cabbibo-favoured decay (CFD) or through mixing followed by doubly-Cabbibo-suppressed decay (DCSD), as shown in Fig. 1. Interference of the two processes introduces the term containing ϕ_3 to the time dependent decay rates, which are given by [5, 6]



(a) CFD (b) DCSD

FIG. 1: Contributions to $B^0 \rightarrow D^{(*)-}\pi^+$ can come either (a) from CFD or (b) from mixing followed by DCSD.

$$\begin{aligned}
P(B^0 \rightarrow D^{(*)+}\pi^-) &= c\,[1 - \cos\Delta m t - 2\Im\bar{\rho}\sin\Delta m t] \\
P(B^0 \rightarrow D^{(*)-}\pi^+) &= c\,[1 + \cos\Delta m t + 2\Im\rho\sin\Delta m t] \\
P(\bar{B}^0 \rightarrow D^{(*)+}\pi^-) &= c\,[1 + \cos\Delta m t + 2\Im\bar{\rho}\sin\Delta m t] \\
P(\bar{B}^0 \rightarrow D^{(*)-}\pi^+) &= c\,[1 - \cos\Delta m t - 2\Im\rho\sin\Delta m t]
\end{aligned} \tag{1}$$

where $c = (e^{-t/\tau_{B^0}})/2\tau_{B^0}$ with τ_{B^0} denoting the lifetime of the neutral B meson and Δm is the B^0-$\bar{B}^0$ mixing parameter. The ρ and $\bar{\rho}$ are defined as $\rho = (q/p)(\mathcal{A}(\bar{B}^0 \rightarrow D^{(*)-}\pi^+)/\mathcal{A}(B^0 \rightarrow D^{(*)-}\pi^+))$ and $\bar{\rho} = (p/q)(\mathcal{A}(B^0 \rightarrow D^{(*)+}\pi^-)/\mathcal{A}(\bar{B}^0 \rightarrow D^{(*)+}\pi^-))$, where p and q relate the mass eigenstates to the flavour eigenstates in the neutral B meson system [5]. They lead to CP violating terms $\Im\rho = -(-1)^L R\sin(2\phi_1+\phi_3-\delta)$ and $\Im\bar{\rho} = (-1)^L R\sin(2\phi_1+\phi_3+\delta)$, where R is the ratio of the magnitudes of the DCSD and CFD amplitudes (here the magnitudes of both the CFD and DCSD amplitudes are assumed to be same for B^0 and $\bar{B}^0$ decays), δ is the strong phase difference between DCSD and CFD, and L is the angular momentum of the final state (1 for $D^*\pi$ and 0 for $D\pi$). R and δ are not necessarily the same for $D^*\pi$ and $D\pi$ final states, and are denoted with subscripts, $D^*\pi$ and $D\pi$, in what follows.

This study uses a $140\,\mathrm{fb}^{-1}$ data sample, which contains 152 million $B\bar{B}$ events, collected with the Belle detector [7] at the KEKB collider [8]. The selection of hadronic events is described elsewhere [9].

For the $\bar{B}^0 \rightarrow D^{*+}\pi^-$ event selection, we use the decay chain $D^{*+} \rightarrow D^0\pi^+$, and $D^0 \rightarrow K^-\pi^+$, $K^-\pi^+\pi^0$ or $K^-\pi^+\pi^+\pi^-$ (charge conjugate modes are implied throughout this paper). For the $\bar{B}^0 \rightarrow D^+\pi^-$ event selection, we use $D^+ \rightarrow K^-\pi^+\pi^+$ decays. Charged tracks except the slow π^+ in the $D^{*+} \rightarrow D^0\pi^+$ decay are required to have a minimum of one hit (two hits) in the r-ϕ (z) plane of the vertex detector in order to allow precise vertex determination. To separate kaons from pions, we form a likelihood for each track, $\mathcal{L}_{K(\pi)}$. The kaon likelihood ratio, $P(K/\pi) = \mathcal{L}_K/(\mathcal{L}_K + \mathcal{L}_\pi)$, has values between 0 (likely to be a pion) and 1 (likely to be a kaon). We require charged kaons to satisfy $P(K/\pi) > 0.3$. No such requirement is imposed to select charged pions coming from D decays.

For D^0 selection, the invariant mass of the daughter particles is required to be within $\pm 16.5\,\mathrm{MeV}/c^2$, $\pm 24.0\,\mathrm{MeV}/c^2$, and $\pm 13.5\,\mathrm{MeV}/c^2$ of the nominal D^0 mass, for $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$ modes, respectively. These intervals correspond to $\pm 3\sigma$, where σ is the Monte Carlo determined invariant mass resolution. For the D^+, the invariant mass is required to be within $\pm 12.5\,\mathrm{MeV}/c^2$ of the nominal D^+ mass. For the $D^0 \to K^-\pi^+\pi^0$ reconstruction, we further require the π^0 momentum to be greater than $200\,\mathrm{MeV}/c$ in the $\Upsilon(4S)$ rest frame, and the ratio of the second to zeroth Fox-Wolfram moments [10] R_2 to be less than 0.55. We require $R_2 < 0.5$ for $D^+ \to K^-\pi^+\pi^+$. We use a mass- and vertex-constrained fit for D^0 and a vertex-constrained fit for D^+.

The D^{*+} is reconstructed by combining D^0 candidates with a slow π^+. Here, slow pions are required to have momentum less than $300\,\mathrm{MeV}/c$ in the $\Upsilon(4S)$ rest frame. The D^* candidates are required to have a mass difference $\Delta M \equiv M_{D^0\pi} - M_{D^0}$ within $\pm 7\,\mathrm{MeV}/c^2$, $\pm 2\,\mathrm{MeV}/c^2$, or $\pm 4\,\mathrm{MeV}/c^2$ of the nominal value, for the $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$ modes respectively.

We reconstruct B candidates by combining the $D^{(*)+}$ candidate with a π^- candidate satisfying $P(K/\pi) < 0.8$. We identify B decays based on requirements on the energy difference $\Delta E \equiv \sum_i E_i - E_{\mathrm{beam}}$ and the beam-energy constrained mass $M_{\mathrm{bc}} \equiv \sqrt{E_{\mathrm{beam}}^2 - (\sum_i \vec{p}_i)^2}$, where E_{beam} is the beam energy, $\vec{p}_i$ and E_i are the momenta and energies of the daughters of the reconstructed B meson candidate, all in the $\Upsilon(4S)$ rest frame. If more than one B candidate is found in the same event, we select the one with best D vertex quality. We define a signal region in the ΔE-M_{bc} plane of $5.27\,\mathrm{GeV}/c^2 < M_{\mathrm{bc}} < 5.29\,\mathrm{GeV}/c^2$ and $|\Delta E| < 0.045\,\mathrm{GeV}$, corresponding to about $\pm 3\sigma$ of both quantities. For the determination of background parameters, we use events in a sideband region defined by $M_{\mathrm{bc}} > 5.2\,\mathrm{GeV}/c^2$ and $-0.14\,\mathrm{GeV} < \Delta E < 0.20\,\mathrm{GeV}$, excluding the signal region.

Charged leptons, pions, and kaons that are not associated with the reconstructed $D^{(*)}\pi$ decays are used to identify the flavour of the accompanying B meson. The algorithm [1] leads to two parameters, q and r, where $q = +1$ indicates $\bar{b}$ hence B^0 and $q = -1$ indicates b hence $\bar{B}^0$. The parameter r is an event-by-event dilution factor ranging from $r = 0$ for no flavour discrimination to $r = 1$ for unambiguous flavour assignment. More than 99.5% of the events are assigned non-zero values of r.

The decay vertices of the $B \to D^{(*)}\pi$ are fitted using the momentum vectors of the D and π (except the slow π from D^* decay) and a requirement that they are consistent with the interaction region profile. For the decay vertices of the tagging B meson, the remaining well reconstructed tracks in the event are used. Tracks that are consistent with K_S^0 decay are rejected. The proper-time difference between the fully reconstructed and the associated B decay is calculated as $\Delta t = (z_{\mathrm{rec}} - z_{\mathrm{tag}})/c\beta\gamma$, where z_{rec} and z_{tag} are the z coordinates of the two B decay vertices and $\beta\gamma = 0.425$ is the Lorentz boost factor at KEKB. After application of the event selection criteria and the requirement that both B's have well defined vertices and $|\Delta t| < 70$ ps ($\sim 45\,\tau_{B^0}$), 7763 and 9351 events remain as the $D^*\pi$ and $D\pi$ candidates, respectively. The signal fractions of the samples, which vary for different r bins, are 96% for $D^*\pi$ and 91% for $D\pi$.

Unbinned maximum likelihood fits to the four time dependent decay rates are performed to extract $\Im\rho$ and $\Im\bar{\rho}$. We minimize $-2\sum_i \ln L_i$ where the likelihood for the i-th event is given by

$$\begin{aligned} L_i &= (1-f_{\rm ol})\left[f_{\rm sig}P_{\rm sig}\otimes R_{\rm sig}+(1-f_{\rm sig})P_{\rm bkg}\otimes R_{\rm bkg}\right] \\ &\quad +f_{\rm ol}P_{\rm ol}. \end{aligned}$$

The signal fraction $f_{\rm sig}$ is determined from the (ΔE, $M_{\rm bc}$) value of each event. The signal distribution is the product of the sum of two Gaussian in ΔE and a Gaussian in $M_{\rm bc}$; that for the background is the product of a first order polynomial in ΔE and an ARGUS function [11] in $M_{\rm bc}$.

The Δt distribution is modeled by a core distribution convolved with resolutions. A small number of events have poorly reconstructed vertices resulting in a very broad distribution Δt. We account for the contributions from these "outliers" by adding a Gaussian component $P_{\rm ol}$ with a width and fraction determined from the B lifetime analysis [12]. The Δt resolution, denoted by $R_{\rm sig}$ and $R_{\rm bkg}$ for the signal and background, is determined on an event-by-event basis, using the estimated uncertainties on the z vertex positions [13].

The signal Δt distributions are given by

$$\begin{aligned} &P_{\rm sig}(q=-1,D^{(*)\pm}\pi^{\mp}) \\ &=(1-w_-)P(B^0\to D^{(*)\pm}\pi^{\mp})+w_+P(\bar{B}^0\to D^{(*)\pm}\pi^{\mp}) \\ &P_{\rm sig}(q=+1,D^{(*)\pm}\pi^{\mp}) \\ &=(1-w_+)P(\bar{B}^0\to D^{(*)\pm}\pi^{\mp})+w_-P(B^0\to D^{(*)\pm}\pi^{\mp}) \end{aligned} \tag{2}$$

where w_- and w_+ are wrong tag fractions for the $q=-1$ and $q=+1$ samples, respectively. P's are given by Eq. 1 with t and c replaced by Δt and $(e^{-|\Delta t|/\tau_{B^0}})/4\tau_{B^0}$, respectively.

The background Δt distribution is parameterized as a sum of a δ-function component and an exponential component with a lifetime $\tau_{\rm bkg}$

$$P_{\rm bkg}=f^{\delta}_{\rm bkg}\delta(\Delta t-\mu^{\delta}_{\rm bkg})+\frac{(1-f^{\delta}_{\rm bkg})}{2\tau_{\rm bkg}}e^{-\left|\Delta t-\mu^{\tau}_{\rm bkg}\right|/\tau_{\rm bkg}} \tag{3}$$

where $f^{\delta}_{\rm bkg}$ is the fraction of events contained in the δ-function, and $\mu^{\delta}_{\rm bkg}$ and $\mu^{\tau}_{\rm bkg}$ are the mean values of $|\Delta t|$ in the δ-function and exponential components, respectively.

While the tagging side should have no asymmetry if the flavour is tagged by primary leptons, it is possible to introduce a small asymmetry when daughter particles of hadronic decays such as $D^{(*)}\pi$ are used for the flavour tagging, due to the same CP violating effect, which is the subject of this paper [14]. This effect is taken into account by replacing the coefficients of $\sin\Delta mt$ in Eqs. 1 by $\Im\bar{\rho}-\Im\bar{\rho}'$, $\Im\rho-\Im\bar{\rho}'$, $\Im\bar{\rho}-\Im\rho'$, and $\Im\rho-\Im\rho'$, respectively. Here the $\Im\rho'$ and $\Im\bar{\rho}'$ represent the CP violating effect due to the presence of $B^0\to\bar{D}X$ and $B^0\to DX$ amplitudes in the flavour tagging side. Note that unlike the $\Im\rho$ and $\Im\bar{\rho}$, which are rigorously defined in terms of $B^0\to D^{(*)\mp}\pi^{\pm}$ and $\bar{B}^0\to D^{(*)\pm}\pi^{\mp}$ amplitudes, $\Im\rho'$ and $\Im\bar{\rho}'$ are effective quantities that include effects of the fraction of $B\to DX$ components in the tagging B decays and all experimental effects of subsequent behaviour of D mesons. Therefore, these quantities must be determined experimentally.

The values of $\Im\rho'$ and $\Im\bar{\rho}'$ are determined in each of six r bins by fitting the Δt distributions of a $D^*l\nu$ control sample [15] using the signal distributions of Eq. 2 and setting $\Im\rho$ and $\Im\bar{\rho}$ to zero. Since the $D^*l\nu$ final states have specific flavour, any observable asymmetry must originate from the tagging side. The results for the combined r bins are $2\Im\rho' = 0.038 \pm 0.014(\mathrm{stat}) \pm 0.005(\mathrm{sys})$ and $2\Im\bar{\rho}' = 0.002 \pm 0.014(\mathrm{stat}) \pm 0.009(\mathrm{sys})$.

The procedures for Δt determination and flavour tagging are tested by extracting τ_{B^0} and Δm. When all four signal categories in Eq. 1 are combined, the signal Δt distribution reduces to an exponential lifetime distribution. We obtain $\tau_{B^0} = 1.583 \pm 0.029$ ps (1.575 ± 0.032 ps) for the $D^*\pi$ ($D\pi$) samples, in good agreement with the world average (1.542 ± 0.016 ps) [3]. Combining the two CFD-dominant modes and the two mixing-dominant modes and ignoring the CP violating terms, the asymmetry behaves as $\cos \Delta m \Delta t$. We obtain $\Delta m = 0.490 \pm 0.015$ ps^{-1} (0.483 ± 0.014 ps^{-1}) for the $D^*\pi$ ($D\pi$) samples, also in good agreement with the world average (0.489 ± 0.008 ps^{-1}) [3]. The same fits also provide wrong tag fractions w_- and w_+ in each r bin for both $D^*\pi$ and $D\pi$ data samples. The errors of our results are statistical only.

We then perform fits to determine the $\Im\rho$ and $\Im\bar{\rho}$ by fixing τ_{B^0} and Δm_d to the world average values and using w_-, w_+, $\Im\rho'$, and $\Im\bar{\rho}'$ for each r bin, as obtained from the above fits. The results are $2\Im\rho_{D^*\pi} = 0.011 \pm 0.057$, $2\Im\bar{\rho}_{D^*\pi} = -0.109 \pm 0.057$, $2\Im\rho_{D\pi} = -0.037 \pm 0.052$, and $2\Im\bar{\rho}_{D\pi} = 0.087 \pm 0.054$. The errors are statistical only. The Δt distributions for the subsamples having the best quality flavour tagging ($0.875 < r < 1.000$) are shown in Fig. 2 for the $D^*\pi$ and in Fig. 3 for the $D\pi$ samples, respectively.

The systematic errors come from i) the uncertainties of parameters which are constrained in the fit, including Δt resolution parameters, background parameters, wrong tag fractions, and physics parameters; ii) uncertainties of the tagging side asymmetries; iii) fit biases induced by the vertexing and other unknown factors. For item i), we repeat the fits varying each parameter value by $\pm 1\sigma$. To estimate item ii), we repeat the fits by varying the $\Im\rho'$ and $\Im\bar{\rho}'$ by their errors. Errors are not explicitly assigned for item iii), since they are included in the errors of $\Im\rho'$ and $\Im\bar{\rho}'$ from the $D^*l\nu$ control sample fit (item ii). Table I summarizes the systematic errors.

TABLE I: Systematic errors in the $2R\sin(2\phi_1 + \phi_3 \pm \delta)$ extractions.

Sources	$D^*\pi$	$D\pi$
Signal Δt resolution	0.014	0.013
Background Δt shape	0.001	0.003
Background fraction	0.002	0.001
Wrong tag fraction	0.006	0.006
Vertexing	0.005	0.005
Physics parameters ($\Delta m, \tau_{B^0}$)	0.001	0.002
Tagging side asymmetry	0.009	0.009
Combined	0.019	0.018

We obtain

$$\begin{aligned}
2R_{D^*\pi}\sin(2\phi_1 + \phi_3 + \delta_{D^*\pi}) &= 0.109 \pm 0.057 \pm 0.019, \\
2R_{D^*\pi}\sin(2\phi_1 + \phi_3 - \delta_{D^*\pi}) &= 0.011 \pm 0.057 \pm 0.019, \\
2R_{D\pi}\sin(2\phi_1 + \phi_3 + \delta_{D\pi}) &= 0.087 \pm 0.054 \pm 0.018, \\
2R_{D\pi}\sin(2\phi_1 + \phi_3 - \delta_{D\pi}) &= 0.037 \pm 0.052 \pm 0.018.
\end{aligned}$$



FIG. 2: Δt distributions for the $D^*\pi$ data in the $0.875 < r < 1.000$ flavour tagging quality bin. (a) $B^0 \to D^{*+}\pi^-$, (b) $B^0 \to D^{*-}\pi^+$, (c) $\bar{B}^0 \to D^{*+}\pi^-$, (d) $\bar{B}^0 \to D^{*-}\pi^+$. Curves show the fit results with the entire event sample, hatched regions indicate the backgrounds.



FIG. 3: Δt distributions for the $D\pi$ events in the $0.875 < r < 1.000$ flavour tagging quality bin. (a) $B^0 \to D^+\pi^-$, (b) $B^0 \to D^-\pi^+$, (c) $\bar{B}^0 \to D^+\pi^-$, (d) $\bar{B}^0 \to D^-\pi^+$. Curves show the fit results with the entire event sample, hatched regions indicate the backgrounds.

The first and second errors are statistical and systematic. At present, the statistical errors are too large to allow any meaningful conclusion to be drawn. However, it is interesting to consider how the four results can be combined using knowledge of R and δ to improve the precision of $\sin(2\phi_1 + \phi_3)$. Several methods have been proposed to measure R [4]. A method that compares the branching fractions of $B^0 \to D_s^{(*)+}\pi^-$ and $B^0 \to D^{(*)-}\pi^+$ and uses factorization relation gives $R_{D\pi} = 0.0237 \pm 0.0050$ and $R_{D^*\pi} = 0.0180 \pm 0.0067$[16]. The present errors are too large to conclude that the two R values are equal. On the other hand, there are solid theoretical grounds for assuming $\delta_{D^*\pi}$ and $\delta_{D\pi}$ to be very small and therefore equal [17]. However, some argue that there is an ambiguity of 180° between $\delta_{D^*\pi}$ and $\delta_{D\pi}$ [6]. Assuming $\delta_{D^{(*)}\pi}$ is close to either 0° or 180°, we obtain $|2R_{D^*\pi}\sin(2\phi_1+\phi_3)| = 0.060 \pm 0.040(\mathrm{stat}) \pm 0.019(\mathrm{sys})$ and $|2R_{D\pi}\sin(2\phi_1+\phi_3)| = 0.061 \pm 0.037(\mathrm{stat}) \pm 0.018(\mathrm{sys})$.

In summary, we measure the time dependent CP violation parameters $2R\sin(2\phi_1+\phi_3\pm\delta)$ for the $B^0(\bar{B}^0) \rightarrow D^{(*)\mp}\pi^{\pm}$ decays using 152 million $B\bar{B}$ events. Under the assumption of $\delta_{D^{(*)}\pi}$ being close to either 0° or 180°, we obtain $|2R_{D^*\pi}\sin(2\phi_1+\phi_3)| = 0.060\pm0.040(\mathrm{stat})\pm 0.019(\mathrm{sys})$ and $|2R_{D\pi}\sin(2\phi_1+\phi_3)| = 0.061\pm0.037(\mathrm{stat})\pm0.018(\mathrm{sys})$.

We wish to thank the KEKB accelerator group for the excellent operation of the KEKB accelerator. We acknowledge support from the Ministry of Education, Culture, Sports, Science, and Technology of Japan and the Japan Society for the Promotion of Science; the Australian Research Council and the Australian Department of Education, Science and Training; the National Science Foundation of China under contract No. 10175071; the Department of Science and Technology of India; the BK21 program of the Ministry of Education of Korea and the CHEP SRC program of the Korea Science and Engineering Foundation; the Polish State Committee for Scientific Research under contract No. 2P03B 01324; the Ministry of Science and Technology of the Russian Federation; the Ministry of Education, Science and Sport of the Republic of Slovenia; the National Science Council and the Ministry of Education of Taiwan; and the U.S. Department of Energy.

---

* on leave from Nova Gorica Polytechnic, Nova Gorica

[1] K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **66** (2002) 071102(R).
[2] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **89** (2002) 201802.
[3] K. Hagiwara *et al.* (Particle Data Group), Phys. Rev. D **66** (2002) 010001.
[4] I. Dunietz and R.G. Sachs, Phys. Rev. D **37** (1988) 3186, *Erratum*: Phys. Rev. D **39** (1989) 3515; I. Dunietz, Phys. Lett. B **427** (1998) 179.
[5] K. Abe, M. Satpathy, and Hitoshi Yamamoto, `hep-ex/0103002`.
[6] R. Fleischer, Nucl. Phys. **B671** (2003) 459.
[7] A. Abashian *et al.* (Belle Collaboration), Nucl. Instr. and Meth. A **479** (2002) 117.
[8] S. Kurokawa, Nucl. Instr. and Meth. A **499** (2003) 1.
[9] K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **66** (2002) 032007.
[10] G.C. Fox and S. Wolfram, Phys. Rev. Lett. **41** (1978) 1581.
[11] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **241** (1990) 278.
[12] H. Tajima, H. Aihara, T. Higuchi *et al.*, `hep-ex/0301026`.
[13] K. Abe, *et al.* (Belle Collaboration), Phys. Rev. Lett. **88** (2002) 171801.
[14] O. Long, M. Baak, R.N. Cahn and D. Kirkby, Phys. Rev. D **68** (2003) 304010.
[15] K. Hara, M. Hazumi *et al.* (Belle Collaboration), Phys. Rev. Lett. **89** (2002) 251803.
[16] We use the PDG values [3] for the $D^{(*)}\pi$ branching fractions, the decay constant ratio $f_{D^{(*)}}/f_{D_s^{(*)}}$ from a recent lattice QCD calculation of K.C. Bowler *et al.*(UKQCD Collaboration), Nucl. Phys. **619** (2001) 507, the $D_s^{(*)}\pi$ branching fractions of P. Krokovny *et al.* (Belle Collaboration), Phys. Rev. Lett. **89** (2002) 231804 and B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **90** (2003) 181803.
[17] L. Wolfenstein, `hep-ph/0309166`.